



**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

SEPTEMBER 2022

TECHNICAL SCIENCES P2

MARKS: 75

TIME: 1½ hours

This question paper consists of 14 pages, including 4 data sheets.

INSTRUCTIONS AND INFORMATION

1. This question paper consists of SEVEN questions. Answer ALL the questions in the ANSWER BOOK.
2. Start EACH question on a NEW page in the ANSWER BOOK.
3. Number the answers correctly according to the numbering system used in this question paper.
4. Leave ONE line between two sub-questions, for example between QUESTION 2.1 and QUESTION 2.2.
5. You may use a non-programmable calculator.
6. You may use appropriate mathematical instruments.
7. You are advised to use the attached DATA SHEETS.
8. Show ALL formulae and substitutions in ALL calculations.
9. Round off your FINAL numerical answers to a minimum of TWO decimal places.
10. Give brief motivations, discussions, etc. where required.
11. Write neatly and legibly.





QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A–D) next to the question numbers (1.1 to 1.5) in the ANSWER BOOK, for example 1.6 D.

1.1. Which ONE of the following general formulae represents alkynes?



1.2 Which ONE of the following compounds represents a ketone?



1.3 Solar cells use p-n junction to convert sunlight directly into a(n) ...

A magnetic field.

B electric field.

C magnetic flux.

D electric current. (2)

1.4 In which ONE of the following options are the three compounds listed in increasing order of vapour pressure?





A propanoic acid, pentane, butan-1-ol

B propanoic acid, butan-1-ol, pentane

C pentane, butan-1-ol, propanoic acid

D butan-1-ol, propanoic acid, pentane (2)

1.5 The cell notation for a standard Zn-Cu electrochemical cell is:



(2)
[10]

QUESTION 2 (Start on a new page.)

Organic chemistry is the chemistry of organic molecules divided into homologous series which are identified by the functional groups.

2.1 Define the term *hydrocarbons*. (2)

2.2 Consider the organic molecules listed below:

A hex-2-ene	B $ \begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & & \text{H} & & \text{O} \\ & & & & & & & & & // \\ \text{H} & - \text{C} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{C} \\ & & & & & & & & & \backslash \\ & \text{H} & & \text{H} & & \text{H} & & \text{H} & & \text{H} \end{array} $
C 3-Chloro-But-1-ene	D $ \begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{O} & & & & \\ & & & & & & & & & \\ \text{H} & - \text{C} & - & \text{C} & - & \text{C} & - & \text{O} & - & \text{H} \\ & & & & & & & & & \\ & \text{H} & & \text{H} & & & & & & \end{array} $


2.2.1 Define the term *isomers* in words. (2)

2.2.2 Draw the structural formula of a positional isomer of **A**. (2)

2.2.3 Write down the name of the homologous series to which **B** belongs. (1)

2.2.4 Give the IUPAC name for the chain isomer of compound **C**. (2)

2.3 The diagram below shows a monomer of the organic compound used for polyethylene. This is the industrial organic product used in the preparation of plastics.



Define the term *monomer* in words. (2)

[11]

QUESTION 3 (Start on a new page.)


The table below shows the vapour pressure of various organic compounds at 25 °C.

Compound	Molar mass (g mol ⁻¹)	Vapour pressure (x10 ² Pa)
pentane	72	573,0
hexane	86	160,0
propan-1-ol	60	21,0
propan-2-ol	60	44,0
butan-1-ol	74	6,2
butan-2-ol	74	18,3
pentan-1-ol	88	2,2
pentan-2-ol	88	8,04
ethanoic acid	60	15,3
propanone	58	240,0

- 3.1 Write down the general formula of the homologous series to which pentane belongs. (1)
- 3.2 Draw the structural formula of propanone. (2)
- 3.3 Give the name of a **functional** isomer of propanone. (1)
- 3.4 Write down the name of the intermolecular forces involved in:
- 3.4.1 Alcohols (1)
- 3.4.2 Alkanes (1)
- 3.5 Refer to the table of organic compounds above to state and explain the relationship between vapour pressure and the strength of intermolecular forces. (2)
- 3.6 Which compound will have the higher boiling point:
Ethanoic acid or propan-1-ol?
- Explain by referring to type of intermolecular forces and energy. (3)
- [11]**

QUESTION 4 (Start on a new page.)

Pent-1-ene can be converted to other compounds by means of different organic reactions represented by **U**, **V**, **W** and **X**, as shown below.



4.1 Write down the TYPE of the reaction represented by:

4.1.1 **U** (1)

4.1.2 **W** (1)

4.1.3 **V** (1)

4.2 During reaction **X**, the alkyl halide (haloalkane) reacts.

4.2.1 Give the NAME of a suitable base used. (1)

4.2.2 Name TWO reaction conditions for reaction **X**. (2)


4.2.3 Write down the balanced reaction using structural formulae for the reaction of Pent-1-ene with hydrogen bromide to form a haloalkane. (3)

4.3 Fossil fuels are formed by the natural process of decomposition of organisms under heat and pressure. They contain a high percentage of carbon and include fuels such as coal, petrol and natural gases. Alkanes are the most important fossil fuels. The combustion of alkanes (also known as oxidation) is highly exothermic.

Write down a balanced reaction for the complete combustion of pentane. (3)
[12]

QUESTION 5 (Start on a new page.)


Semiconductor devices such as diodes are widely used in modern electronics.



- 5.1 Define the term *semiconductor* in words. (2)
- 5.2 Phosphorus was added to silicon in small quantities. It was then found that the electrical conductivity of silicon improved.
- 5.2.1 Identify the described process in the above statement. (1)
- 5.2.2 What type of a semiconductor material (P-type or N-type) is formed during this process? Give a reason for your answer. (2)
- [5]**

QUESTION 6 (Start on a new page.)

In the electrolytic cell, represented below, two CARBON RODS are used as electrodes and a concentrated copper (II) chloride solution is used as an electrolyte.



When the cell is functioning, the following **observations** are made:

- A gas is released at electrode **P**
- Electrode **Q** is coated with a reddish-brown layer

6.1 Define the term *electrolyte*. (2)

6.2 Write down a half-reaction to explain the observation made at:

6.2.1 Electrode **P** (2)

6.2.2 Electrode **Q** (2)

6.3 Write down the energy conversion that is taking place in this cell. (1)

6.4 Which electrode, **P** or **Q**, is the cathode? Give a reason for your answer. (2)

6.5 The carbon rods in the above cell are now replaced with COPPER RODS. The following observations are made at electrode **P**:

- No gas is released
- Its surface appears rough and eroded

6.5.1 Refer to the RELATIVE STRENGTHS OF REDUCING AGENTS to explain this observation. (3)


6.5.2 This cell can be used for the electroplating of a bracelet in the industry. Which electrode (**P** or **Q**) will be replaced with a bracelet during the electroplating process? (1)

[13]

QUESTION 7 (Start on a new page.)

The potential difference of a galvanic cell, measured experimentally by learners in a Technical Sciences laboratory, is COMPARED with its potential difference calculated at standard conditions.

They set up the galvanic cell shown below.



The voltmeter measures an initial reading of **0,19 V**.

7.1 Write down the energy conversion that takes place in this cell. (1)

7.2 State ONE function of the salt bridge. (1)

7.3 Write down the half-reaction that takes place at the anode. (2)

7.4 In which direction do electrons flow in the external circuit when this cell delivers a current?

Write down only **FROM Sn TO Cd** or **FROM Cd TO Sn**. (1)

7.5 Write down the balanced net (overall) cell reaction. (3)

7.6 Use the Table of STANDARD REDUCTION POTENTIALS to calculate the initial potential difference (emf) of the above cell at STANDARD CONDITIONS. (3)

7.7 From the results obtained, the learners concluded that the measured potential difference differs from the calculated potential difference.

Give TWO possible reasons for this difference in values. (2)

[13]

TOTAL: 75

**NATIONAL SENIOR CERTIFICATE
NASIONALE SENIOR SERTIFIKAAT**

**DATA FOR TECHNICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR TEGNIESE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)**

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESE KONSTANTES

NAAM/NAME	SIMBOOL/SYMBOL	WAARDE/VALUE
Standard pressure <i>Standaarddruk</i>	p^θ	$1,013 \times 10^5 \text{ Pa}$
Standard temperature <i>Standaardtemperatuur</i>	T^θ	273 K

TABLE 2: FORMULAE/TABEL 2: FORMULES

$E^\theta_{\text{cell}} = E^\theta_{\text{cathode}} - E^\theta_{\text{anode}} / E^\theta_{\text{sel}} = E^\theta_{\text{katode}} - E^\theta_{\text{anode}}$ $E^\theta_{\text{cell}} = E^\theta_{\text{reduction}} - E^\theta_{\text{oxidation}} / E^\theta_{\text{sel}} = E^\theta_{\text{reduksie}} - E^\theta_{\text{oksidasie}}$ $E^\theta_{\text{cell}} = E^\theta_{\text{oxidising agent}} - E^\theta_{\text{reducing agent}} / E^\theta_{\text{sel}} = E^\theta_{\text{oksideermiddel}} - E^\theta_{\text{reduseermiddel}}$

TABLE 3: THE PERIODIC TABLE OF ELEMENTS/TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE

1 (I)	2 (II)	3	4	5	6	7	8	9	10	11	12	13 (III)	14 (IV)	15 (V)	16 (VI)	17 (VII)	18 (VIII)
<p>KEY/ SLEUTEL</p> <p><i>Atoomgetal</i> Atomic number</p> <p><i>Elektronegatiwiteit</i> Electronegativity</p> <p><i>Simbool</i> Symbol</p> <p><i>Benaderde relatiewe atoommassa</i> Approximate relative atomic mass</p>																	
1 H 1																	2 He 4
3 Li 7	4 Be 9											5 B 11	6 C 12	7 N 14	8 O 16	9 F 19	10 Ne 20
11 Na 23	12 Mg 24											13 Al 27	14 Si 28	15 P 31	16 S 32	17 Cl 35,5	18 Ar 40
19 K 39	20 Ca 40	21 Sc 45	22 Ti 48	23 V 51	24 Cr 52	25 Mn 55	26 Fe 56	27 Co 59	28 Ni 59	29 Cu 63,5	30 Zn 65	31 Ga 70	32 Ge 73	33 As 75	34 Se 79	35 Br 80	36 Kr 84
37 Rb 86	38 Sr 88	39 Y 89	40 Zr 91	41 Nb 92	42 Mo 96	43 Tc 98	44 Ru 101	45 Rh 103	46 Pd 106	47 Ag 108	48 Cd 112	49 In 115	50 Sn 119	51 Sb 122	52 Te 128	53 I 127	54 Xe 131
55 Cs 133	56 Ba 137	57 La 139	72 Hf 179	73 Ta 181	74 W 184	75 Re 186	76 Os 190	77 Ir 192	78 Pt 195	79 Au 197	80 Hg 201	81 Tl 204	82 Pb 207	83 Bi 209	84 Po 209	85 At 210	86 Rn 222
87 Fr 223	88 Ra 226	89 Ac 227															
			58 Ce 140	59 Pr 141	60 Nd 144	61 Pm 147	62 Sm 150	63 Eu 152	64 Gd 157	65 Tb 159	66 Dy 163	67 Ho 165	68 Er 167	69 Tm 169	70 Yb 173	71 Lu 175	
			90 Th 232	91 Pa 231	92 U 238	93 Np 237	94 Pu 244	95 Am 243	96 Cm 247	97 Bk 247	98 Cf 251	99 Es 252	100 Fm 257	101 Md 288	102 No 289	103 Lr 260	

TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD REDUKSIEPOTENSIALE

Increasing oxidising ability/Toenemende oksiderende vermoë

Half-reactions/Halfreaksies			E^{θ} (V)
$F_2(g) + 2e^-$	\rightleftharpoons	$2F^-$	+ 2,87
$Co^{3+} + e^-$	\rightleftharpoons	Co^{2+}	+ 1,81
$H_2O_2 + 2H^+ + 2e^-$	\rightleftharpoons	$2H_2O$	+ 1,77
$MnO_4^- + 8H^+ + 5e^-$	\rightleftharpoons	$Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^-$	\rightleftharpoons	$2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^-$	\rightleftharpoons	$2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^-$	\rightleftharpoons	$2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^-$	\rightleftharpoons	$Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^-$	\rightleftharpoons	Pt	+ 1,20
$Br_2(l) + 2e^-$	\rightleftharpoons	$2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^-$	\rightleftharpoons	$NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^-$	\rightleftharpoons	$Hg(l)$	+ 0,85
$Ag^+ + e^-$	\rightleftharpoons	Ag	+ 0,80
$NO_3^- + 2H^+ + e^-$	\rightleftharpoons	$NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^-$	\rightleftharpoons	Fe^{2+}	+ 0,77
$O_2(g) + 2H^+ + 2e^-$	\rightleftharpoons	H_2O_2	+ 0,68
$I_2 + 2e^-$	\rightleftharpoons	$2I^-$	+ 0,54
$Cu^+ + e^-$	\rightleftharpoons	Cu	+ 0,52
$SO_2 + 4H^+ + 4e^-$	\rightleftharpoons	$S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^-$	\rightleftharpoons	$4OH^-$	+ 0,40
$Cu^{2+} + 2e^-$	\rightleftharpoons	Cu	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^-$	\rightleftharpoons	$SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^-$	\rightleftharpoons	Cu^+	+ 0,16
$Sn^{4+} + 2e^-$	\rightleftharpoons	Sn^{2+}	+ 0,15
$S + 2H^+ + 2e^-$	\rightleftharpoons	$H_2S(g)$	+ 0,14
$2H^+ + 2e^-$	\rightleftharpoons	$H_2(g)$	0,00
$Fe^{3+} + 3e^-$	\rightleftharpoons	Fe	- 0,06
$Pb^{2+} + 2e^-$	\rightleftharpoons	Pb	- 0,13
$Sn^{2+} + 2e^-$	\rightleftharpoons	Sn	- 0,14
$Ni^{2+} + 2e^-$	\rightleftharpoons	Ni	- 0,27
$Co^{2+} + 2e^-$	\rightleftharpoons	Co	- 0,28
$Cd^{2+} + 2e^-$	\rightleftharpoons	Cd	- 0,40
$Cr^{3+} + e^-$	\rightleftharpoons	Cr^{2+}	- 0,41
$Fe^{2+} + 2e^-$	\rightleftharpoons	Fe	- 0,44
$Cr^{3+} + 3e^-$	\rightleftharpoons	Cr	- 0,74
$Zn^{2+} + 2e^-$	\rightleftharpoons	Zn	- 0,76
$2H_2O + 2e^-$	\rightleftharpoons	$H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^-$	\rightleftharpoons	Cr	- 0,91
$Mn^{2+} + 2e^-$	\rightleftharpoons	Mn	- 1,18
$Al^{3+} + 3e^-$	\rightleftharpoons	Al	- 1,66
$Mg^{2+} + 2e^-$	\rightleftharpoons	Mg	- 2,36
$Na^+ + e^-$	\rightleftharpoons	Na	- 2,71
$Ca^{2+} + 2e^-$	\rightleftharpoons	Ca	- 2,87
$Sr^{2+} + 2e^-$	\rightleftharpoons	Sr	- 2,89
$Ba^{2+} + 2e^-$	\rightleftharpoons	Ba	- 2,90
$Cs^+ + e^-$	\rightleftharpoons	Cs	- 2,92
$K^+ + e^-$	\rightleftharpoons	K	- 2,93
$Li^+ + e^-$	\rightleftharpoons	Li	- 3,05

Increasing reducing ability/Toenemende reduserende vermoë

TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD REDUKSIEPOTENSIALE

Half-reactions/Halfreaksies			E^{θ} (V)
$\text{Li}^+ + \text{e}^-$	\rightleftharpoons	Li	-3,05
$\text{K}^+ + \text{e}^-$	\rightleftharpoons	K	-2,93
$\text{Cs}^+ + \text{e}^-$	\rightleftharpoons	Cs	-2,92
$\text{Ba}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ba	-2,90
$\text{Sr}^{2+} + 2\text{e}^-$	\rightleftharpoons	Sr	-2,89
$\text{Ca}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ca	-2,87
$\text{Na}^+ + \text{e}^-$	\rightleftharpoons	Na	-2,71
$\text{Mg}^{2+} + 2\text{e}^-$	\rightleftharpoons	Mg	-2,36
$\text{Al}^{3+} + 3\text{e}^-$	\rightleftharpoons	Al	-1,66
$\text{Mn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Mn	-1,18
$\text{Cr}^{2+} + 2\text{e}^-$	\rightleftharpoons	Cr	-0,91
$2\text{H}_2\text{O} + 2\text{e}^-$	\rightleftharpoons	$\text{H}_2(\text{g}) + 2\text{OH}^-$	-0,83
$\text{Zn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Zn	-0,76
$\text{Cr}^{3+} + 3\text{e}^-$	\rightleftharpoons	Cr	-0,74
$\text{Fe}^{2+} + 2\text{e}^-$	\rightleftharpoons	Fe	-0,44
$\text{Cr}^{3+} + \text{e}^-$	\rightleftharpoons	Cr^{2+}	-0,41
$\text{Cd}^{2+} + 2\text{e}^-$	\rightleftharpoons	Cd	-0,40
$\text{Co}^{2+} + 2\text{e}^-$	\rightleftharpoons	Co	-0,28
$\text{Ni}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ni	-0,27
$\text{Sn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Sn	-0,14
$\text{Pb}^{2+} + 2\text{e}^-$	\rightleftharpoons	Pb	-0,13
$\text{Fe}^{3+} + 3\text{e}^-$	\rightleftharpoons	Fe	-0,06
$2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{H}_2(\text{g})$	0,00
$\text{S} + 2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{H}_2\text{S}(\text{g})$	+0,14
$\text{Sn}^{4+} + 2\text{e}^-$	\rightleftharpoons	Sn^{2+}	+0,15
$\text{Cu}^{2+} + \text{e}^-$	\rightleftharpoons	Cu^+	+0,16
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+0,17
$\text{Cu}^{2+} + 2\text{e}^-$	\rightleftharpoons	Cu	+0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4\text{e}^-$	\rightleftharpoons	4OH^-	+0,40
$\text{SO}_2 + 4\text{H}^+ + 4\text{e}^-$	\rightleftharpoons	$\text{S} + 2\text{H}_2\text{O}$	+0,45
$\text{Cu}^+ + \text{e}^-$	\rightleftharpoons	Cu	+0,52
$\text{I}_2 + 2\text{e}^-$	\rightleftharpoons	2I^-	+0,54
$\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	H_2O_2	+0,68
$\text{Fe}^{3+} + \text{e}^-$	\rightleftharpoons	Fe^{2+}	+0,77
$\text{NO}_3^- + 2\text{H}^+ + \text{e}^-$	\rightleftharpoons	$\text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+0,80
$\text{Ag}^+ + \text{e}^-$	\rightleftharpoons	Ag	+0,80
$\text{Hg}^{2+} + 2\text{e}^-$	\rightleftharpoons	$\text{Hg}(\ell)$	+0,85
$\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^-$	\rightleftharpoons	$\text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0,96
$\text{Br}_2(\ell) + 2\text{e}^-$	\rightleftharpoons	2Br^-	+1,07
$\text{Pt}^{2+} + 2\text{e}^-$	\rightleftharpoons	Pt	+1,20
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1,23
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^-$	\rightleftharpoons	$2\text{H}_2\text{O}$	+1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^-$	\rightleftharpoons	$2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1,33
$\text{Cl}_2(\text{g}) + 2\text{e}^-$	\rightleftharpoons	2Cl^-	+1,36
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^-$	\rightleftharpoons	$\text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1,51
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$2\text{H}_2\text{O}$	+1,77
$\text{Co}^{3+} + \text{e}^-$	\rightleftharpoons	Co^{2+}	+1,81
$\text{F}_2(\text{g}) + 2\text{e}^-$	\rightleftharpoons	2F^-	+2,87

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reduserende vermoë